

Molecular packing, piezo- and pyroelectric properties of *tert*-butyl *N*-(*tert*-butoxycarbonyl)-(*S*)-prolinamide

D.A. Gruzdev^{1,2}, A.S. Nuraeva³, S.G. Vasilev³, A.V. Ushakov³, V.O. Ustinova¹,
P.A. Slepukhin^{1,2}, V.P. Krasnov^{1,2}, V.Ya. Shur³

¹Postovsky Institute of Organic Synthesis, Russian Academy of Sciences (Ural Branch), 620990, Ekaterinburg, Russia

e-mail: gruzdev-da@ios.uran.ru

²Institute of Chemical Engineering, Ural Federal University, 620002, Ekaterinburg, Russia

³School of Natural Sciences and Mathematics, Ural Federal University, 620000, Ekaterinburg, Russia

Chiral organic compounds attract considerable attention as a basis for biocompatible and environmental friendly materials. It is known that natural amino acid derivatives are capable of self-organization in solid state [1] and possess piezoelectric [2] and pyroelectric properties [3].

We studied the properties of crystals of *N*-(*tert*-butoxycarbonyl)-(*S*)-prolinamide **1**. Compound **1** was obtained as a result of coupling between *N*-Boc-(*S*)-proline and *tert*-butylamine by the method of mixed anhydrides. The single crystals of **1** were grown by spontaneous crystallization from a water–methanol mixture and by a vapor diffusion technique.

It has been found that compound **1** forms crystals of two types differing in their morphology. X-Ray diffraction analysis (XDA) has shown that the crystals belong to hexagonal (*hex*-**1**, space group *P*6₁) or triclinic syngony (*tc*-**1**, space group *P*1) (Fig. 1). The general motif of molecular packing of *hex*-**1** is a formation of nanotubes oriented along the main crystal axis.




Figure 1. The structural formula and molecular packing of compound **1** according to XDA.

The magnitudes of vertical piezoelectric response of isomorphous crystals *hex*-**1** measured independently by piezoresponse force microscopy and interferometry techniques were in good agreement. The observed piezocoefficients reached 12 pC/N.

The pyroelectric effect of crystals *hex*-**1** was measured.

It has been shown that the phase transition is induced by the IR laser irradiation. The single crystals *hex*-**1** transformed to polymorphic *hex*-**1** / *tc*-**1** conglomerates.

The work was financially supported by the Russian Foundation for Basic Research (grant no. 16-33-60122). The equipment of the Ural Centre for Shared Use “Modern Nanotechnologies” UrFU and Centre for Joint Use “Spectroscopy and Analysis of Organic Compounds” was used.

1. L. Adler-Abramovich, E. Gazit, *Chem. Soc. Rev.* **43**, 6881 (2014).
2. S. Vasilev, P. Zelenovskiy, D. Vasileva, A. Nuraeva, V.Ya. Shur, A.L. Kholkin, *J. Phys. Chem. Solids* **93**, 68 (2016).
3. S. Piperno, E. Mirzadeh, E. Mishuk, D. Ehre, S. Cohen, M. Eisenstein, M. Lahav, I. Lubomirsky, *Angew. Chem. Int. Ed.* **52**, 6513 (2013).